

Stochastic Two-Stage Steiner Tree

Thomas Kesselheim

Last Update: November 30, 2025

This time, we will consider another example of stochastic two-stage optimization. Last time, we considered LP-based approaches. One of the drawbacks of these approaches is that one always has to solve a linear program whose size depends on the number of scenarios. Today, we will consider a different kind of algorithm that is more combinatorial. The number of scenarios does not matter at all. Indeed, we only need to be able to draw samples from the same distribution that the scenario is generated from.

1 Recap: Steiner Tree

We consider a stochastic variant of the rooted Steiner tree problem. Recall that in the deterministic offline problem we are given a graph $G = (V, E)$, edge weights $w_e \geq 0$ for $e \in E$, a root $r \in V$, and a set of terminals $T \subseteq V$. Our task is to select a subset of the edges $S \subseteq E$ such that $\{r\} \cup T$ is connected in $G' = (V, S)$ and $\sum_{e \in S} w_e$ is minimized. It is an NP-hard problem. Observe that if $T = V$ then this problem is exactly the minimum spanning tree problem. Without loss of generality, $G = (V, E)$ is a complete graph. We can also assume that the weights w_e fulfill the triangle inequality. That is, $w_{\{u,v\}} \leq w_{\{u,x\}} + w_{\{x,v\}}$ for all $u, v, x \in V$. This is without loss of generality because we could instead take the detour via x instead of the edge $\{u, v\}$.

Also recall that Steiner trees can be approximated by minimum spanning trees. Such a spanning tree only uses edges between the nodes in the set $\{r\} \cup T$ and no edges to other vertices (called Steiner vertices). Let $\text{MST}(T) \subseteq E$ be the minimum spanning tree on $G|_{\{r\} \cup T}$ and let $\text{Steiner}(T) \subseteq E$ be the optimal Steiner tree connecting $\{r\} \cup T$. We already proved the following lemma when we talked about Online Steiner Tree.

Lemma 15.1. *A minimum spanning tree on $G|_{\{r\} \cup T}$ is a 2-approximation for the min-cost Steiner tree on $\{r\} \cup T$, formally*

$$w(\text{MST}(T)) \leq 2 \cdot w(\text{Steiner}(T))$$

2 Two-Stage Problem

In the stochastic variant, we do not know the set T in advance but only the distribution it is drawn from. As a matter of fact, we do not need to fully know the probability distribution that T is drawn from. It will only be necessary to be able to draw samples from the same distribution.

In the first stage, we do not yet know the set T but we can already pick edges e at costs w_e . In the second stage, we know the set T but edges are more expensive now: Picking edge e costs $\lambda \cdot w_e$ for $\lambda \geq 1$. That is, we assume that cost of edges increase by a uniform inflation factor $\lambda \geq 1$ from the first stage to the second.

Our goal is to minimize the expected cost. The expected cost of a policy is

$$\sum_{e \text{ selected in first stage}} w_e + \mathbf{E} \left[\sum_{e \text{ selected in second stage}} \lambda \cdot w_e \right].$$

Let us understand the limiting cases first: In the case $\lambda = 1$ it does not make sense to select anything in the first stage because it does not get more expensive in the second one. For $\lambda \rightarrow \infty$, the second stage gets extremely expensive, so we buy edges connecting every possible T in the first stage.

Again, even the basic Steiner tree problem is NP hard. Therefore, we cannot compute the optimal policy in polynomial time and we want to approximate it instead. More formally, let E_0^* be the set of edges selected by the optimal policy in the first stage, and let E_T^* be the set of edges selected by the optimal policy in the second stage if the set of terminals is T . We are looking for a policy whose expected cost is as close as possible to

$$Z^* := \sum_{e \in E_0^*} w_e + \mathbf{E} \left[\sum_{e \in E_T^*} \lambda \cdot w_e \right].$$

3 Algorithm “Boosted Sampling”

For simplicity, we will assume that λ is an integer. We will consider the following algorithm called “Boosted Sampling”:

- In the first stage, draw λ times from the known distribution, call these sets $S_1, \dots, S_\lambda$. Compute a minimum spanning tree on $\{r\} \cup S_1 \cup \dots \cup S_\lambda$, let E_0 be the set of edges contained in it and pick them.
- In the second stage, set $w_e = 0$ for all $e \in E_0$ and compute a minimum spanning tree on $\{r\} \cup T$, let E_T be the set of contained edges not picked so far and pick them.

This algorithm only needs to sample λ times and calculate two minimum spanning trees. It therefore runs in polynomial time if λ is polynomially bounded.

Theorem 15.2. *The expected cost of the algorithm is at most $4Z^*$. That is,*

$$\mathbf{E} \left[\sum_{e \in E_0} w_e + \sum_{e \in E_T} \lambda \cdot w_e \right] \leq 4Z^*.$$

In the proof, we will bound the expected cost from each stage by $2Z^*$. By linearity of expectation, this then implies

$$\mathbf{E} \left[\sum_{e \in E_0} w_e + \sum_{e \in E_T} \lambda \cdot w_e \right] = \mathbf{E} \left[\sum_{e \in E_0} w_e \right] + \mathbf{E} \left[\sum_{e \in E_T} \lambda \cdot w_e \right] \leq 2Z^* + 2Z^* = 4Z^*.$$

4 Analysis of First Stage

Lemma 15.3. *The expected first-stage cost of the algorithm is at most $2Z^*$. That is,*

$$\mathbf{E} \left[\sum_{e \in E_0} w_e \right] \leq 2Z^*.$$

Proof. By our notation, $E_0^* \cup E_{S_1}^* \cup \dots \cup E_{S_\lambda}^*$ is the union of all edges chosen by the optimal policy for scenarios $S_1, \dots, S_\lambda$. Observe that $E_0^* \cup E_{S_1}^* \cup \dots \cup E_{S_\lambda}^*$ is a feasible Steiner tree connecting all of $S_1 \cup \dots \cup S_\lambda$ to the root r .

Our choice, $E_0 = \text{MST}(S_1 \cup \dots \cup S_\lambda)$ can have at most twice the cost, so

$$w(E_0) \leq 2w(E_0^* \cup E_{S_1}^* \cup \dots \cup E_{S_\lambda}^*) \leq 2w(E_0^*) + 2 \sum_{i=1}^{\lambda} w(E_{S_i}^*) .$$

By linearity of expectation, we have

$$\mathbf{E} [w(E_0)] \leq 2w(E_0^*) + 2 \sum_{i=1}^{\lambda} \mathbf{E} [w(E_{S_i}^*)] .$$

Furthermore, observe that $\mathbf{E} [w(E_{S_i}^*)] = \mathbf{E} [w(E_T^*)]$ for all i because S_i and T are drawn from the same distribution. So

$$\mathbf{E} [w(E_0)] \leq 2w(E_0^*) + 2\lambda \mathbf{E} [w(E_T^*)] = 2Z^* . \quad \square$$

5 Analysis of Second Stage

Lemma 15.4. *The expected second-stage cost of the algorithm is at most $2Z^*$. That is,*

$$\mathbf{E} \left[\sum_{e \in E_T} \lambda \cdot w_e \right] \leq 2Z^* .$$

To bound the cost incurred in the second stage, we have to understand how expensive it is to “augment” a spanning tree. Given $A, B \subseteq V$, let $\delta(A, B)$ be the cost of a minimum spanning tree on the graph $G|_{\{r\} \cup A \cup B}$ when setting $w_{\{u,v\}} = 0$ for all $u, v \in \{r\} \cup A$. That is, the vertices in $\{r\} \cup A$ can be connected for free but we also have to connect all of B to them.

Lemma 15.5. *For any $U_1, \dots, U_k \subseteq V$, we have*

$$\sum_{i=1}^k \delta \left(\bigcup_{j \neq i} U_j, U_i \right) \leq w(\text{MST}(U_1 \cup \dots \cup U_k)) .$$




Figure 1: Illustration of Lemma 15.5 with two sets U_1 and U_2 . Using only the red edges, each red vertex is connected to the root or a blue vertex, which we can connect for free, or is blue itself. The same holds if we swap red and blue.

Proof. Consider $\text{MST}(U_1 \cup \dots \cup U_k)$. Recall that this is a tree rooted at r . For $v \in U_1 \cup \dots \cup U_k$, $v \neq r$, let a_v be the weight of the edge connecting v to its parent node in this tree.

Now, we can bound

$$\delta \left(\bigcup_{j \neq i} U_j, U_i \right) \leq \sum_{v \in U_i \setminus \bigcup_{j \neq i} U_j} a_v$$

because by connecting each $v \in U_i \setminus \bigcup_{j \neq i} U_j$ to its parent node and using the zero-weight edges all of $U_1 \cup \dots \cup U_k$ is connected (also see Figure 1).

Therefore, we now have

$$\sum_{i=1}^k \delta \left(\bigcup_{j \neq i} U_j, U_i \right) \leq \sum_{i=1}^k \sum_{v \in U_i \setminus \bigcup_{j \neq i} U_j} a_v \leq \sum_{v \in \bigcup_i U_i} a_v = w(\text{MST}(U_1 \cup \dots \cup U_k)) . \quad \square$$

Based on this lemma, we can now complete the analysis of the second stage.

Proof of Lemma 15.4. In the second stage, we connect the set T by augmenting a minimum spanning tree on $\{r\} \cup S_1 \cup \dots \cup S_\lambda$ to one that also includes the set T . Therefore

$$\sum_{e \in E_T} \lambda \cdot w_e = \lambda \cdot \delta(S_1 \cup \dots \cup S_\lambda, T) .$$

We now perform a thought experiment: Note that $S_1, \dots, S_\lambda$ and T are $\lambda + 1$ independent draws from the same distribution. So, equivalently, we might also draw $U_1, \dots, U_{\lambda+1}$ from this distribution and then draw K uniformly at random from $\{1, \dots, \lambda + 1\}$ and set $T = U_K$ and assign the other U_i sets arbitrarily to $S_1, \dots, S_\lambda$.

Therefore, we can write

$$\mathbf{E} [\delta(S_1 \cup \dots \cup S_\lambda, T)] = \mathbf{E} \left[\delta \left(\bigcup_{j \neq K} U_j, U_K \right) \right] = \mathbf{E} \left[\frac{1}{\lambda + 1} \sum_{i=1}^{\lambda+1} \delta \left(\bigcup_{j \neq i} U_j, U_i \right) \right] .$$

By Lemma 15.5, we have

$$\sum_{i=1}^{\lambda+1} \delta \left(\bigcup_{j \neq i} U_j, U_i \right) \leq w(\text{MST}(U_1 \cup \dots \cup U_{\lambda+1})) .$$

So, combining these arguments, the second-stage cost of our algorithm can be bounded by

$$\mathbf{E} \left[\sum_{e \in E_T} \lambda \cdot w_e \right] \leq \frac{\lambda}{\lambda + 1} \mathbf{E} [w(\text{MST}(U_1 \cup \dots \cup U_{\lambda+1}))] .$$

Again, $E_0^* \cup E_{U_1}^* \cup \dots \cup E_{U_{\lambda+1}}^*$ is a feasible Steiner tree connecting $U_1 \cup \dots \cup U_{\lambda+1}$ to the root, so the minimum spanning tree can have at most twice the cost, formally

$$\begin{aligned} w(\text{MST}(U_1 \cup \dots \cup U_{\lambda+1})) &\leq 2w(E_0^* \cup E_{U_1}^* \cup \dots \cup E_{U_{\lambda+1}}^*) \\ &\leq 2w(E_0^*) + 2 \sum_{i=1}^{\lambda+1} w(E_{U_i}^*) . \end{aligned}$$

Again use linearity of expectation and $\mathbf{E} [w(E_{U_i}^*)] = \mathbf{E} [w(E_T^*)]$ to get

$$\mathbf{E} [w(\text{MST}(U_1 \cup \dots \cup U_{\lambda+1}))] \leq 2w(E_0^*) + 2(\lambda + 1) \mathbf{E} [w(E_T^*)] \leq 2 \frac{\lambda + 1}{\lambda} (w(E_0^*) + \lambda \mathbf{E} [w(E_T^*)]) . \quad \square$$

Reference

Boosted sampling: approximation algorithms for stochastic optimization, A. Gupta, M. Pál, R. Ravi, A. Sinha, STOC 2004