

## Basics of Online Convex Optimization, Part 1

Thomas Kesselheim

Last Update: January 18, 2026

Today, we will get to know a much larger framework for online learning. Indeed, the experts setting will come back as a special case and also the multiplicative-weights algorithm. As a motivating example, we will consider linear regression. In its simplest case, one is given a number of pairs  $(x^{(t)}, y^{(t)})$  of data points. One then computes a line, defined by a slope  $w_1$  and a  $y$ -intercept  $w_2$  so as to minimize the squared error  $\sum_t (w_1 x^{(t)} + w_2 - y^{(t)})^2$ . One can then use this line to predict the  $y$ -label given an  $x$ -coordinate.




Figure 1: The red line is the regression line, which minimized the sum of the squared errors.

We turn this problem into an online problem as follows. We will see the data points  $(x^{(1)}, y^{(1)}), (x^{(2)}, y^{(2)}), \dots$  one after the other. Indeed, we will first only see  $x^{(t)}$  and have to predict  $y^{(t)}$  before we get to know its actual value. That is, we already have to predict while learning.

One could also picture a kind of “bandit” feedback for this model: Instead of getting to know the actual  $y^{(t)}$ , we only get to know how far we are off the actual value.

## 1 General Setup

We consider the following round-based problem. We will have to optimize a sequence of a priori unknown functions  $f_1, \dots, f_T$ . Each  $f_t$  maps from set  $S$  to the real numbers. The set  $S \subseteq \mathbb{R}^d$  is a set of  $d$ -dimensional real vectors.

In each step  $t \in \{1, \dots, T\}$ , we choose a point  $\mathbf{w}^{(t)} \in S$ . Only afterwards, we get to know  $f_t$  and incur a cost of  $f_t(\mathbf{w}^{(t)})$ .

The *regret* of a sequence  $\mathbf{w}^{(1)}, \dots, \mathbf{w}^{(T)}$  is defined as before as the amount by which our decisions are more expensive than the best single point in hindsight. That is,

$$\text{Regret}^{(T)} = \sum_{t=1}^T f_t(\mathbf{w}^{(t)}) - \min_{\mathbf{u} \in S} \sum_{t=1}^T f_t(\mathbf{u}) .$$

Note that, as in the last lectures, the order of minimum and sum in the second term is important.

**Example 22.1.** To capture our example of simple linear regression, we can set  $d = 2$  and  $S = \mathbb{R}^2$ . A point  $(w_1, w_2) \in S$  corresponds to the slope  $w_1$  and the  $y$ -intercept  $w_2$  of the regression line. A function  $f_t$  is the square of the error that we make on the  $t$ -th sample, depending on which  $w_1$  and  $w_2$  we use. So

$$f_t(w_1, w_2) = \left( w_1 x^{(t)} + w_2 - y^{(t)} \right)^2 .$$

Note that the best single  $(w_1, w_2)$  in hindsight corresponds exactly to the optimal regression line.

If the set  $S$  is finite, we could run the algorithm for experts and bandits problems. In our regression example it is infinite. Instead we will assume that  $S$  and the functions  $f_1, \dots, f_T$  are convex.

## 2 Convex Sets, Convex Functions, and Gradients

We assume that each function  $f_t$  is differentiable<sup>1</sup> and convex.

The typical example one should keep in mind is a quadratic function in one dimension (see Figure 2). One way to define convexity in this setting is to require that the function never falls below its tangents. This is expressed in terms of the derivative as follows. For all  $u, v \in S$  we have to have

$$f(u) \geq f(v) + f'(v)(u - v) .$$




Figure 2: A typical convex function in one dimension, including a tangent in red.

In multiple dimensions, the idea is just the same. The function  $f$  now has a gradient  $\nabla f$ , which is defined to be the vector of all partial derivatives;  $(\nabla f(\mathbf{u}))_i = \frac{\partial f}{\partial u_i}(\mathbf{u})$ . A function  $f$  is convex if it never falls below the tangent hyperplane (see Figure 3). That is for all  $\mathbf{u}, \mathbf{v}$

$$f(\mathbf{u}) \geq f(\mathbf{v}) + \langle \nabla f(\mathbf{v}), (\mathbf{u} - \mathbf{v}) \rangle . \quad (1)$$

Here  $\langle \cdot, \cdot \rangle$  denotes the inner product, defined by  $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^d x_i y_i$ .

**Example 22.2.** Another important—and familiar—example is the following. Set  $S = \{ \mathbf{v} \in \mathbb{R}^d \mid v_i \geq 0 \text{ for all } i, \sum_{i=1}^d v_i = 1 \}$ . The functions  $f_t$  are linear. That is,  $f_t(\mathbf{v}) = \sum_{i=1}^d \ell_i^{(t)} v_i$  for some  $\ell_i^{(t)} \in \mathbb{R}$ . These functions are clearly convex. And we already know this setting: It's the experts setting with  $d$  experts and the vectors  $\mathbf{v} \in S$  correspond to probability distributions over experts.

<sup>1</sup>None of the results actually requires differentiability but the exposition gets a lot easier.




Figure 3: A convex function in two dimensions.

### 3 Follow-the-Leader

A very natural algorithm is the following *Follow the Leader*: In every step  $t$ , choose the point  $\mathbf{w}^{(t)}$  that would have resulted in the cheapest cost up to now, that is, set  $\mathbf{w}^{(t)}$  to  $\mathbf{v}$  such that  $\sum_{t'=1}^{t-1} f_{t'}(\mathbf{v})$  is minimal. The point  $\mathbf{w}^{(1)}$  is arbitrary.

What we would actually want to do is to also include the function  $f_t$  in the sum because this determines the actual cost in step  $t$ . Unfortunately, we do not know it when choosing  $\mathbf{w}^{(t)}$  but only when choosing  $\mathbf{w}^{(t+1)}$ . Our first observation is that we can bound the regret by the distances of  $\mathbf{w}^{(t)}$  and  $\mathbf{w}^{(t+1)}$ .

**Lemma 22.3.** *For Follow-the-Leader, we have*

$$\text{Regret}^{(T)} \leq \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{w}^{(t+1)})) .$$

*Proof.* We have to show that for all  $T \geq 0$

$$\sum_{t=1}^T f_t(\mathbf{w}^{(t)}) - \min_{\mathbf{u} \in S} \sum_{t=1}^T f_t(\mathbf{u}) \leq \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{w}^{(t+1)})) ,$$

or equivalently

$$\min_{\mathbf{u} \in S} \sum_{t=1}^T f_t(\mathbf{u}) \geq \sum_{t=1}^T f_t(\mathbf{w}^{(t+1)}) .$$

We will show this bound by induction on  $T$ .

The statement is trivial for  $T = 0$ . For  $T > 0$ , we may assume by induction hypothesis that it already holds for  $T - 1$ . So, we get

$$\sum_{t=1}^{T-1} f_t(\mathbf{w}^{(T+1)}) \geq \min_{\mathbf{u} \in S} \sum_{t=1}^{T-1} f_t(\mathbf{u}) \stackrel{\text{ind. hyp.}}{\geq} \sum_{t=1}^{T-1} f_t(\mathbf{w}^{(t+1)}) .$$

By adding  $f_T(\mathbf{w}^{(T+1)})$  to both sides, we get

$$\sum_{t=1}^T f_t(\mathbf{w}^{(T+1)}) \geq \sum_{t=1}^T f_t(\mathbf{w}^{(t+1)}) .$$

Recall the definition of  $\mathbf{w}^{(T+1)}$ . It is chosen such that  $\sum_{t=1}^T f_t(\mathbf{w}^{(T+1)})$  is minimized, which means nothing but

$$\min_{\mathbf{u} \in S} \sum_{t=1}^T f_t(\mathbf{u}) = \sum_{t=1}^T f_t(\mathbf{w}^{(T+1)}) .$$

In combination, these two bounds show the claim for  $T$ . □

**Example 22.4.** Let  $S = [-1, 1]$  and

$$f_1(w) = \frac{w}{2} \quad f_{2k}(w) = -w \quad f_{2k+1}(w) = w \quad \text{for all } k \in \mathbb{N} .$$

In odd steps  $t \geq 3$ ,  $\sum_{t'=1}^{t-1} f_{t'}(w) = -\frac{w}{2}$ ; in even steps  $t$ ,  $\sum_{t'=1}^{t-1} f_{t'}(w) = \frac{w}{2}$ . Therefore, Follow-the-leader chooses  $w^{(1)}$  arbitrarily,  $w^{(2)} = -1$ ,  $w^{(3)} = 1$ ,  $w^{(4)} = -1$ ,  $\dots$ . Therefore  $f_1(w^{(1)}) \geq -\frac{1}{2}$  and  $f_t(w^{(t)}) = 1$  for all  $t > 1$ . Choosing, in contrast,  $u = 0$ , then for all  $t$  we get  $f_t(u) = 0$ . So,  $\text{Regret}^{(T)} \geq T - \frac{3}{2}$ .

## 4 Follow-the-Regularized-Leader

The problem in Example 22.4 is that the optimal point keeps jumping from one extreme to the other; Follow-the-Leader is always “too late”. Therefore, we modify the algorithm a tiny bit. We add a *regularization term*: We choose  $\mathbf{w}^{(t)}$  as the  $\mathbf{v}$  that minimizes  $R(\mathbf{v}) + \sum_{t'=1}^{t-1} f_{t'}(\mathbf{v})$ . The function  $R: S \rightarrow \mathbb{R}$  is a suitable function that has higher values for more “extreme” values.

**Example 22.5.** Typical choices of regularizers are

- Euclidean regularization

$$R(\mathbf{v}) = \frac{1}{2\eta} \sum_{i=1}^d v_i^2 ,$$

- Entropical regularization (for non-negative vectors)

$$R(\mathbf{v}) = \frac{1}{\eta} \sum_{i=1}^d v_i \ln v_i ,$$

where  $\eta > 0$  is a scaling factor, determining how strong the regularization works. Smaller values of  $\eta$  mean stronger regularization. In the case of Euclidean regularization, points closer to the origin are preferred. Entropical regularization prefers values between 0 and 1 to the boundary points.

Recall that  $S = \{\mathbf{v} \in \mathbb{R}^d \mid v_i \geq 0 \text{ for all } i, \sum_{i=1}^d v_i = 1\}$  with linear functions  $f_i$  is exactly the experts setting. One can show that Entropical regularization makes us choose  $\mathbf{w}^{(t)}$  exactly such that  $w_i^{(t)}$  is proportional to  $\exp(-\eta \sum_{t'=1}^{t-1} \ell_i^{(t')})$ . This is exactly the multiplicative-weights update rule.

**Example 22.6.** Let us see what happens in Example 22.4 with Euclidean regularization. In odd steps  $t \geq 3$ ,  $\sum_{t'=1}^{t-1} f_{t'}(w) + R(w) = -\frac{w}{2} + \frac{1}{2\eta} w^2$ ; in even steps  $t$ ,  $\sum_{t'=1}^{t-1} f_{t'}(w) + R(w) = \frac{w}{2} + \frac{1}{2\eta} w^2$ . These are minimized by  $w^{(t)} = \frac{\eta}{2}$  for odd  $t$  and  $w^{(t)} = -\frac{\eta}{2}$  for even  $t$ . So, if  $\eta$  is small enough, we indeed keep close to the origin.

We can extend the regret bound from above to Follow-the-Regularized-Leader.

**Lemma 22.7.** *For Follow-the-Regularized-Leader, we have*

$$\text{Regret}^{(T)} \leq \max_{\mathbf{u} \in S} R(\mathbf{u}) - R(\mathbf{w}^{(1)}) + \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{w}^{(t+1)})) .$$

*Proof.* Follow-the-Regularized-Leader is nothing but Follow-the-Leader with a hypothetical “step zero”, in which  $f_0 = R$ . So, Lemma 22.3 tells us that for all  $\mathbf{u} \in S$

$$\sum_{t=0}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{u})) \leq \sum_{t=0}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{w}^{(t+1)})) ,$$

which now means

$$R(\mathbf{w}^{(0)}) - R(\mathbf{u}) + \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{u})) \leq R(\mathbf{w}^{(0)}) - R(\mathbf{w}^{(1)}) + \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{w}^{(t+1)})) .$$

Because this bound holds for all  $\mathbf{u} \in S$ , rearranging gives us

$$\text{Regret}^{(T)} = \max_{\mathbf{u} \in S} \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{u})) \leq \max_{\mathbf{u} \in S} R(\mathbf{u}) - R(\mathbf{w}^{(1)}) + \sum_{t=1}^T (f_t(\mathbf{w}^{(t)}) - f_t(\mathbf{w}^{(t+1)})) . \quad \square$$

At first sight, this regret bound might look weaker than the one for Follow-the-Leader. The point is, however, that the regularization keeps the difference of  $f_t(\mathbf{w}^{(t)})$  and  $f_t(\mathbf{w}^{(t+1)})$  smaller if it is chosen in a suitable way.